

SYNOPSIS

01/31/2020

Review of “Novel coronavirus 2019-nCoV: early estimation of epidemiological parameters and epidemic predictions”

Article citation: Read JM, Bridgen JRE, Cummings DAT, Ho A, Jewell CP. Novel coronavirus 2019-nCoV: early estimation of epidemiological parameters and epidemic predictions. medRxiv. 2020 Jan 28 [Epub ahead of print]. Available from: <https://www.medrxiv.org/content/10.1101/2020.01.23.20018549v2>

Note: This is a synopsis of the version of this article posted on January 28, 2020 and may be subject to change.

One-minute summary

- This study used case information from Chinese cities and other countries from January 1-22, 2020 to fit a mathematical model to estimate outbreak parameters
- The model suggests that:
 - **early estimate of the basic reproductive number (R0) was 3.1 (95%CI 2.4-4.1)**, which is higher both than SARS and MERS, but similar to some early SARS estimates
 - **58-76% of transmissions must be prevented** by control measures for infections to stop increasing
 - **Only 5.0% (95%CI 3.6-7.4) of infections in Wuhan are identified**, indicating that the number of infections in Wuhan is large and the majority of infections may be mild and not detected
- Without change to transmission or control, the model suggests that:
 - On January 29, 2020, there will be 105,077 infected individuals (prediction interval: 46,635-185,412) in Wuhan
 - Other **cities in China with largest predicted outbreaks** include: Shanghai, Beijing, Guangzhou, Shenzhen, Chengdu, and Kunming
 - Jurisdictions with the **greatest predicted risk of importing infections through air travel** include: Thailand, Japan, South Korea, Taiwan, Hong Kong, USA, Singapore, Malaysia, Australia, and Vietnam

Additional information

- Model based on several assumptions and is specific to time period and data available
- Findings are critically dependent on underlying assumptions:
 - No control interventions factored in
 - Variables driving epidemic dynamics are constant
 - Travel behaviour continues as per mobility estimates
 - Only air travel considered; land transport not considered
 - Assumed latent period of 4 days
 - Considered only human-to-human transmission

PHO reviewer's comments

- Model predictions are based on data from January 1-22, 2020 and a number of assumptions that have not been validated; predictions may change as further data become available
- Article not peer reviewed

Citation

Ontario Agency for Health Protection and Promotion (Public Health Ontario). Review of “Novel coronavirus 2019-nCoV: early estimation of epidemiological parameters and epidemic predictions”. Toronto, ON: Queens’s Printer for Ontario; 2020.

Disclaimer

This document was developed by Public Health Ontario (PHO). PHO provides scientific and technical advice to Ontario’s government, public health organizations and health care providers. PHO’s work is guided by the current best available evidence at the time of publication.

The application and use of this document is the responsibility of the user. PHO assumes no liability resulting from any such application or use.

This document may be reproduced without permission for non-commercial purposes only and provided that appropriate credit is given to PHO. No changes and/or modifications may be made to this document without express written permission from PHO.

Public Health Ontario

Public Health Ontario is a Crown corporation dedicated to protecting and promoting the health of all Ontarians and reducing inequities in health. Public Health Ontario links public health practitioners, front-line health workers and researchers to the best scientific intelligence and knowledge from around the world. For more information about PHO, visit publichealthontario.ca.

