

WEEKLY EPIDEMIOLOGICAL SUMMARY

SARS-CoV-2 Genomic Surveillance in Ontario, February 3, 2025

This report summarizes the results of SARS-CoV-2 whole genome sequencing completed by the Ontario COVID-19 Genomics Network as of January 29, 2025.

Background

The continued monitoring of global SARS-CoV-2 genomic data has identified changes in the virus' genome as it spreads through populations. These random changes or mutations arise as a virus evolves over time. The accumulation of these mutations can result in a new lineage of the virus, which is a common occurrence. These new lineages will differ slightly in genome sequence and are termed variants. Although many variants will have no difference in the ability to spread or cause disease, some variants have mutations which may enhance virulence, transmissibility, and/or allow the virus to escape natural or vaccine-induced immunity.

The identification of variants and mutations occurs through whole genome sequencing (WGS) of select samples. Through global surveillance of SARS-CoV-2 genomes, a number of variants have been identified with evidence of clinical and/or public health significance, termed variants of concern (VOC). Variants designated as VOCs include B.1.1.7 (Alpha), B.1.351 (Beta), P.1 (Gamma), B.1.617.2 (Delta), and B.1.1.529 (Omicron), some of which have been de-escalated due to their diminishing prevalence globally.¹⁻³

As SARS-CoV-2 continues to evolve, lineages will naturally divide into descendant sublineages - a genetically closely related group derived from a common ancestor. The descendant branches are given new lineage aliases, such as for B.1.1.529 (Omicron) lineages (e.g., BA.2 [alias for B.1.1.529.2] and JN.1 [alias for B.1.1.529.2.86.1.1]). When a host is infected with two or more descendant lineages, lineages can recombine to form a new recombinant lineage (e.g., KP.3.3 and KS.1.1 to form XEC). New designations represent refined genetic groups that can be tracked separately. As more research is conducted, there may be evidence of an important difference in terms of transmissibility, severity, or immune escape, at which time WHO may assign a new Greek letter classification to a lineage.

The Ontario COVID-19 Genomics Network (OCGN) performs WGS on all eligible positive SARS-CoV-2 samples (see Technical Notes for details). Sequences are processed using bioinformatics analyses and assigned a Pango lineage⁴ using the pangolin tool⁵, allowing for the identification of lineages.

Highlights

- In the most recent week (January 12 to January 18), a total of 693 cases were sequenced. XEC was the most prevalent lineage (30.7%), followed by KP.3.1.1 (12.7%), and LP.8.1 (10.0%).
- The proportion of XEC decreased from 32.6% (January 5 to January 11) to 30.7% (January 12 to January 18).
 - Based on the Nowcast model, XEC is projected to decrease to 27.0% (95% CI: 22.9% - 31.7%) by February 5, 2025.
- The proportion of KP.3.1.1 remained stable at 11.9% (January 5 to January 11) and 12.7% (January 12 to January 18).
 - Based on the Nowcast model, KP.3.1.1 is projected to decrease to 6.5% (95% CI: 5.0% - 8.3%) by February 5, 2025. The weekly relative growth rate of KP.3.1.1 is 0.87 (95% CI: 0.83 - 0.92) times that of XEC.
- The proportion of LP.8.1 increased from 8.8% (January 5 to January 11) to 10.0% (January 12 to January 18).
 - Based on the Nowcast model, LP.8.1 is projected to increase to 22.5% (95% CI: 16.7% - 29.8%) by February 5, 2025. The weekly relative growth rate of LP.8.1 is 1.35 (95% CI: 1.25 - 1.45) times that of XEC.

Lineage counts and designations may change between reports as components of the Pango lineage assignment models are updated (see Technical Notes for details).

Due to technical and logistical issues, data from Kingston Health Sciences Centre are incomplete for week 52, 2024 to week 3, 2025 (December 22, 2024 to January 18, 2025), data from Hamilton Regional Laboratory Medicine Program is incomplete for week 3 (January 12 to January 18), and data from The Hospital for Sick Children are incomplete for week 2 to week 3 (January 5 to January 18). As a result, counts, proportions, and Nowcast projections may change compared to previous and subsequent reports.

Representative Surveillance


Table 1: Number of SARS-CoV-2 positive specimens, number and percentage of specimens sequenced for representative surveillance by week, Ontario, December 22, 2024 to January 18, 2025

Week	Number of Positive Specimens	Number of Specimens Sequenced	Percentage of Specimens Sequenced
Week 52 (December 22 - December 28)	1,866	1,125	60.3%
Week 1 (December 29 - January 4)	2,311	1,260	54.5%
Week 2 (January 5 - January 11)	1,990	1,006	50.6%
Week 3 (January 12 - January 18)	1,655	703	42.5%
Total	7,822	4,094	52.3%

Note: The 4,094 specimens sequenced were associated with 3,904 unique cases; in the most recent week, 703 specimens sequenced were associated with 693 unique cases. Unique cases are the denominator for tables throughout the report. ‘Number of positive specimens’ is the number of tests positive for SARS-CoV-2 in Ontario. Date was assigned to best align with sample collection date, which may differ from other PHO products. ‘Number of specimens sequenced’ is the number of specimens sequenced for representative surveillance. ‘Percentage sequenced’ may be lower than the sampling proportion because not all specimens are eligible to be sequenced (i.e. excludes samples with cycle threshold >30 or insufficient volume). Results may not be representative of Ontario overall. For representative surveillance: details on the proportion of eligible samples sequenced by the OCGN can be found in the Technical Notes. Week was assigned based on earliest date available for a sample. Not all sequencing and bioinformatics analyses for the most recent weeks were complete at the time of data extraction. Case counts for these weeks may increase in subsequent reports.

Data sources: Ontario Laboratories Information System (OLIS) from the Ontario Respiratory Virus Tool (ORVT), Ontario Health Data Platform - Public Health Analytic Environment (OHDP-PHAE)

Figure 1: Percentage of SARS-CoV-2 cases by the most prevalent lineages and week, representative surveillance, Ontario, January 21, 2024 to January 18, 2025



Note: Results may not be representative of Ontario overall. Details on the proportion of eligible samples sequenced by the OCGN can be found in the Technical Notes. Week was assigned based on earliest date available for a sample. If more than one sample was sequenced for a case, the most recent sample was included. Not all sequencing and bioinformatics analyses for the most recent weeks were complete at the time of data extraction. Case counts for these weeks may increase in subsequent reports.

Data sources: Ontario Health Data Platform - Public Health Analytic Environment (OHDP-PHAE)


Table 2: Number and percentage of SARS-CoV-2 cases by Pango lineage and week, representative surveillance, Ontario, December 22, 2024 to January 18, 2025

Pango Lineage	Week 52 (December 22 - December 28)	Week 1 (December 29 - January 4)	Week 2 (January 5 - January 11)	Week 3 (January 12 - January 18)	Total (December 22 - January 18)
XEC	332 (30.6%)	427 (36.7%)	314 (32.6%)	213 (30.7%)	1,286 (32.9%)
KP.3.1.1	195 (18.0%)	174 (15.0%)	115 (11.9%)	88 (12.7%)	572 (14.7%)
MC.1	113 (10.4%)	83 (7.1%)	90 (9.3%)	62 (8.9%)	348 (8.9%)
LP.8.1	60 (5.5%)	74 (6.4%)	85 (8.8%)	69 (10.0%)	288 (7.4%)
MC.10.1	45 (4.1%)	72 (6.2%)	52 (5.4%)	53 (7.6%)	222 (5.7%)
XEC.2	38 (3.5%)	42 (3.6%)	52 (5.4%)	17 (2.5%)	149 (3.8%)
NP.1	44 (4.1%)	44 (3.8%)	38 (3.9%)	12 (1.7%)	138 (3.5%)
XEK	38 (3.5%)	30 (2.6%)	23 (2.4%)	29 (4.2%)	120 (3.1%)
XEC.4	14 (1.3%)	18 (1.5%)	24 (2.5%)	13 (1.9%)	69 (1.8%)
MC.13	13 (1.2%)	29 (2.5%)	14 (1.5%)	6 (0.9%)	62 (1.6%)
Other recombinant	27 (2.5%)	25 (2.2%)	19 (2.0%)	20 (2.9%)	91 (2.3%)
Other Omicron	166 (15.3%)	144 (12.4%)	138 (14.3%)	111 (16.0%)	559 (14.3%)
Total sequenced	1,085 (100%)	1,162 (100%)	964 (100%)	693 (100%)	3,904 (100%)

Note: Includes the most prevalent lineages detected in the past month. Details on the proportion of eligible samples sequenced by the OCGN can be found in the Technical Notes. Week was assigned based on the earliest date available for the sample. Not all sequencing and bioinformatics analyses for the most recent weeks were complete at the time of data extraction. Case counts for these weeks may increase in subsequent reports.

Data sources: Ontario Health Data Platform - Public Health Analytic Environment (OHDP-PHAE)

Figure 2: Estimated daily SARS-CoV-2 prevalence (%) by Pango lineage, using Nowcast model, Ontario, December 8, 2024 to February 8, 2025



Note: Each curve represents the estimated prevalence of a given lineage from Nowcast modelling, which uses six weeks of daily representative surveillance data in a multinomial logistic regression. Each set of dots represents the observed daily prevalence of a given lineage, while their size represents the relative number of samples. The vertical dashed line indicates the most recent day of data, after which projected Nowcast prevalence estimates are presented with their 95% confidence intervals. The vertical grey lines indicate the mid-point of the week. Lineages with at least 21 days of non-zero case counts were included in the model and lineages that did not have at least 21 days of non-zero case counts were included but not shown. Figure includes all lineages with at least one day of an estimated prevalence of 5% or greater during the 12 week period (six observed and six projected). Only three weeks of projected data are shown. Prevalence projections may be overestimated for emerging lineages.

Data sources: Ontario Health Data Platform - Public Health Analytic Environment (OHDP-PHAE)

Table 3: Estimated SARS-CoV-2 prevalence (%), projected prevalence (%), and weekly relative growth rate (with 95% confidence intervals) by Pango lineage, using Nowcast model, Ontario, December 8, 2024 to February 8, 2025

Pango Lineage	Week 2 (January 8): Estimated	Week 3 (January 15): Estimated	Week 4 (January 22): Projected	Week 5 (January 29): Projected	Week 6 (February 5): Projected	Weekly Relative Growth Rate
XEC	32.9 (31.2 - 34.6)	32.4 (30.1 - 34.7)	31.2 (28.3 - 34.3)	29.4 (25.9 - 33.2)	27.0 (22.9 - 31.7)	1.00 (reference)
KP.3.1.1	13.6 (12.4 - 14.8)	11.6 (10.3 - 13.2)	9.8 (8.3 - 11.5)	8.0 (6.5 - 9.9)	6.5 (5.0 - 8.3)	0.87 (0.83 - 0.92)
MC.1	8.9 (7.9 - 10.0)	8.5 (7.3 - 10.0)	8.0 (6.5 - 9.8)	7.4 (5.7 - 9.5)	6.6 (4.8 - 9.0)	0.98 (0.92 - 1.04)
LP.8.1	8.4 (7.4 - 9.4)	11.1 (9.5 - 13.0)	14.4 (11.7 - 17.6)	18.2 (14.1 - 23.2)	22.5 (16.7 - 29.8)	1.35 (1.25 - 1.45)
MC.10.1	6.3 (5.5 - 7.3)	7.4 (6.1 - 8.9)	8.4 (6.6 - 10.7)	9.4 (6.9 - 12.7)	10.2 (7.0 - 14.6)	1.18 (1.09 - 1.28)
NP.1	3.1 (2.5 - 3.8)	2.6 (2.0 - 3.4)	2.1 (1.5 - 2.9)	1.6 (1.1 - 2.5)	1.3 (0.8 - 2.1)	0.84 (0.77 - 0.91)
XEC.4	2.1 (1.6 - 2.7)	2.8 (2.0 - 3.9)	3.7 (2.3 - 5.7)	4.7 (2.7 - 8.1)	5.9 (3.0 - 11.3)	1.36 (1.18 - 1.58)
Other lineages	9.9 (8.9 - 11.0)	8.5 (7.3 - 9.9)	7.2 (5.9 - 8.7)	5.9 (4.7 - 7.5)	4.8 (3.6 - 6.4)	0.88 (0.83 - 0.92)

Note: The Nowcast model uses six weeks of daily representative surveillance data in a multinomial logistic regression that estimates and projects the prevalence of SARS-CoV-2 lineages. The weekly relative growth rate is a measure of a lineage’s growth rate relative to the reference lineage and is estimated in the Nowcast model. The weekly relative growth rate and projections may be overestimated for emerging lineages. The prevalence estimates and projections presented are from the Wednesday (mid-point) of the specified week. Lineages with at least 21 days of non-zero case counts were included in the model separately. ‘Other lineages’ includes all other lineages combined that did not individually have at least 21 days of non-zero case counts. Lineages that had at least one day with a prevalence of 5% or greater in the 12 week period (six observed and six projected) were included in the table. Only two weeks of observed and three weeks of projected data are shown. Prevalence estimates are based on the model and are not expected to be the same as the observed data (e.g. Table 2). Details on the methodology used to calculate Nowcast prevalence estimates, projections, and the weekly relative growth rates can be found in the Technical Notes.

Data sources: Ontario Health Data Platform - Public Health Analytic Environment (OHDP-PHAE)

Table 4: Number and percentage of SARS-CoV-2 cases by Pango lineage and age group, representative surveillance, Ontario, December 22, 2024 to January 18, 2025

Pango Lineage	Ages: 0-4	Ages: 5-11	Ages: 12-19	Ages: 20-39	Ages: 40-59	Ages: 60-79	Ages: 80 and over	Total
XEC	57 (36.5%)	4 (30.8%)	6 (33.3%)	59 (30.4%)	105 (33.5%)	358 (31.8%)	696 (33.5%)	1,285 (32.9%)
KP.3.1.1	25 (16.0%)	1 (7.7%)	0 (0.0%)	29 (14.9%)	37 (11.8%)	191 (17.0%)	288 (13.8%)	571 (14.6%)
MC.1	7 (4.5%)	1 (7.7%)	2 (11.1%)	10 (5.2%)	26 (8.3%)	108 (9.6%)	193 (9.3%)	347 (8.9%)
LP.8.1	12 (7.7%)	0 (0.0%)	4 (22.2%)	23 (11.9%)	35 (11.2%)	71 (6.3%)	142 (6.8%)	287 (7.4%)
MC.10.1	4 (2.6%)	1 (7.7%)	1 (5.6%)	9 (4.6%)	9 (2.9%)	55 (4.9%)	143 (6.9%)	222 (5.7%)
XEC.2	7 (4.5%)	1 (7.7%)	0 (0.0%)	8 (4.1%)	6 (1.9%)	44 (3.9%)	83 (4.0%)	149 (3.8%)
NP.1	9 (5.8%)	1 (7.7%)	1 (5.6%)	12 (6.2%)	11 (3.5%)	47 (4.2%)	57 (2.7%)	138 (3.5%)
XEK	2 (1.3%)	0 (0.0%)	0 (0.0%)	8 (4.1%)	13 (4.2%)	35 (3.1%)	62 (3.0%)	120 (3.1%)
XEC.4	2 (1.3%)	0 (0.0%)	0 (0.0%)	2 (1.0%)	4 (1.3%)	16 (1.4%)	45 (2.2%)	69 (1.8%)
MC.13	3 (1.9%)	1 (7.7%)	0 (0.0%)	3 (1.5%)	4 (1.3%)	19 (1.7%)	32 (1.5%)	62 (1.6%)
Other recombinant	5 (3.2%)	0 (0.0%)	0 (0.0%)	6 (3.1%)	10 (3.2%)	25 (2.2%)	45 (2.2%)	91 (2.3%)
Other Omicron	23 (14.7%)	3 (23.1%)	4 (22.2%)	25 (12.9%)	53 (16.9%)	157 (13.9%)	294 (14.1%)	559 (14.3%)
Total sequenced	156 (100%)	13 (100%)	18 (100%)	194 (100%)	313 (100%)	1,126 (100%)	2,080 (100%)	3,900 (100%)

Note: Includes the most prevalent lineages detected in the past month. Age was assigned based on the birth date provided in OCGN; excludes cases with missing birth dates.

Data sources: Ontario Health Data Platform - Public Health Analytic Environment (OHDP-PHAE)

Table 5: Number and percentage of SARS-CoV-2 cases by Pango lineage and geographic region, representative surveillance, Ontario, December 22, 2024 to January 18, 2025

Pango Lineage	North West	North East	Eastern	Central East	Toronto	South West	Central West	Unknown	Total
XEC	17 (33.3%)	15 (16.9%)	84 (28.3%)	335 (30.3%)	195 (27.5%)	170 (42.1%)	258 (39.6%)	212 (35.4%)	1,286 (32.9%)
KP.3.1.1	10 (19.6%)	4 (4.5%)	35 (11.8%)	157 (14.2%)	121 (17.1%)	72 (17.8%)	68 (10.4%)	105 (17.5%)	572 (14.7%)
MC.1	1 (2.0%)	10 (11.2%)	36 (12.1%)	107 (9.7%)	59 (8.3%)	31 (7.7%)	59 (9.0%)	45 (7.5%)	348 (8.9%)
LP.8.1	0 (0.0%)	4 (4.5%)	25 (8.4%)	83 (7.5%)	79 (11.2%)	13 (3.2%)	43 (6.6%)	41 (6.8%)	288 (7.4%)
MC.10.1	0 (0.0%)	1 (1.1%)	8 (2.7%)	86 (7.8%)	38 (5.4%)	24 (5.9%)	24 (3.7%)	41 (6.8%)	222 (5.7%)
XEC.2	0 (0.0%)	1 (1.1%)	10 (3.4%)	47 (4.3%)	44 (6.2%)	14 (3.5%)	18 (2.8%)	15 (2.5%)	149 (3.8%)
NP.1	16 (31.4%)	2 (2.2%)	8 (2.7%)	43 (3.9%)	34 (4.8%)	3 (0.7%)	14 (2.1%)	18 (3.0%)	138 (3.5%)
XEK	0 (0.0%)	2 (2.2%)	35 (11.8%)	17 (1.5%)	5 (0.7%)	12 (3.0%)	29 (4.4%)	20 (3.3%)	120 (3.1%)
XEC.4	0 (0.0%)	12 (13.5%)	1 (0.3%)	4 (0.4%)	5 (0.7%)	6 (1.5%)	29 (4.4%)	12 (2.0%)	69 (1.8%)
MC.13	0 (0.0%)	0 (0.0%)	10 (3.4%)	17 (1.5%)	2 (0.3%)	8 (2.0%)	21 (3.2%)	4 (0.7%)	62 (1.6%)
Other recombinant	1 (2.0%)	0 (0.0%)	9 (3.0%)	27 (2.4%)	14 (2.0%)	10 (2.5%)	21 (3.2%)	9 (1.5%)	91 (2.3%)
Other Omicron	6 (11.8%)	38 (42.7%)	36 (12.1%)	181 (16.4%)	112 (15.8%)	41 (10.1%)	68 (10.4%)	77 (12.9%)	559 (14.3%)
Total sequenced	51 (100%)	89 (100%)	297 (100%)	1,104 (100%)	708 (100%)	404 (100%)	652 (100%)	599 (100%)	3,904 (100%)

Note: Cases with missing/unassigned patient postal code (15.2%) or out of province postal codes (0.1%) were included in the “Unknown” category. Sample date represents the earliest date available for the sample. Not all sequencing and bioinformatics analyses for the most recent weeks were complete at the time of data extraction. Case counts for these weeks may increase in subsequent reports. Geographic region was assigned based on OCGN patient postal code.

Data sources: Ontario Health Data Platform - Public Health Analytic Environment (OHDP-PHAE)

Cumulative Whole Genome Sequencing Results

Table 6: Number of SARS-CoV-2 cases by Pango lineage, cumulative counts, Ontario, December 22, 2024 to January 18, 2025

WHO Label / Pango Lineage	December 22, 2024 - January 18, 2025
Omicron	2,189
JN.1.16	1
JN.1.16.5	4
JN.1.18.6	1
KP.2.14	1
KP.2.3.4	28
KP.2.6	1
KP.3	1
KP.3.1	2
KP.3.1.1	572
KP.3.2.3	2
KP.3.2.6	3
KP.3.3	28
KP.3.3.1	5
KP.3.3.2	32
KP.3.3.4	2
KP.3.7	1
KP.4.2	9
KS.1.1	1
LB.1.3.1	32
LF.7	8
LF.7.1	7
LF.7.1.2	4
LF.7.2	1
LF.7.2.1	2
LF.7.3	12
LP.7	11
LP.8.1	288
MA.1	1
MB.1	1
MC.1	348
MC.1.1	6

WHO Label / Pango Lineage	December 22, 2024 - January 18, 2025
MC.10	19
MC.10.1	222
MC.10.2	49
MC.11	13
MC.13	62
MC.13.1	4
MC.13.2	4
MC.14	2
MC.16	10
MC.17	2
MC.19	58
MC.2	16
MC.21	2
MC.21.1	12
MC.22	7
MC.4	2
MC.6	21
MC.8	21
MC.8.1	53
MC.9	2
NC.1	24
NF.1	3
NL.1	1
NL.2	1
NL.3	14
NL.4	9
NL.5	3
NP.1	138
Recombinant	1,715
NB.1	1
XDY	1
XDY.1	2
XEC	1,286
XEC.1	3
XEC.2	149
XEC.3	7
XEC.4	69

WHO Label / Pango Lineage	December 22, 2024 - January 18, 2025
XEC.5	20
XEC.6	3
XEF	26
XEK	120
XEL	19
XEM	9
Total Sequenced	4,187

Note: Results do not represent all Ontario cases. Includes results from the OHDP-PHAE from the past month. Pango lineage assignments may change over time, which may impact cumulative totals. Results should be interpreted with caution as frequencies do not reflect prevalence. Sample date represents the earliest date available for the sample. If more than one sample was sequenced for a case, the most recent sample was included.

Data sources: Ontario Health Data Platform - Public Health Analytic Environment (OHDP-PHAE)

Technical Notes

Data Sources

Ontario Health Data Platform – Public Health Analytic Environment (OHDP-PHAE)

- Ontario COVID-19 Genomics Network (OCGN) data were extracted from the Ontario Health Data Platform – Public Health Analytic Environment on January 30, 2025 at approximately 9:00 a.m.

Public Health Ontario (PHO)

- Data were submitted to the OHDP-PHAE on January 28, 2025 at approximately 11:00 a.m.

The Hospital for Sick Children (HSC)

- Data were submitted to the OHDP-PHAE on January 28, 2025 at approximately 10:45 a.m.

Kingston Health Sciences Centre (KHSC)

- Data were submitted to the OHDP-PHAE on December 10, 2024 at approximately 1:45 p.m.

Shared Hospital Laboratory (SHL)

- Data were submitted to the OHDP-PHAE on January 28, 2025 at approximately 10:30 a.m.

Hamilton Regional Laboratory Medicine Program (HRLMP)

- Data were submitted to the OHDP-PHAE on January 22, 2025 at approximately 1:30 p.m.

Ontario Laboratories Information System (OLIS) data – Ontario Respiratory Virus Tool (ORVT)

- OLIS data were extracted from Public Health Ontario’s ORVT on January 31, 2025 at approximately 11:30 a.m.

Ontario SARS-CoV-2 Whole Genome Sequencing Strategy

- Ontario’s whole genome sequencing strategy began early 2021 to confirm the identification of VOCs from PCR testing. Since then, the strategy has shifted to representative surveillance as of May 2, 2021. Diagnostic testing laboratories currently send all eligible samples (diagnostic PCR Ct \leq 30 and sufficient volume remaining) to one of the five OCGN laboratories for whole genome sequencing.
- The Ministry of Health continues to update its [guidance on testing](#) and as such, representative surveillance only pertains to tested populations.

Data Caveats and Methods: Ontario COVID-19 Genomics Network (OCGN)

- Lineage is assigned using the Phylogenetic Assignment of Named Global Outbreak Lineages (pangolin) tool, a software package for predicting SARS-CoV-2 lineages from genome sequences and global lineages. Lineages were reported using pangolin version 4.3.1, pangolin data version 1.31, pangolin assignment version 1.31, scorpio version 0.3.19, and constellations version 0.1.12.
- Lineage nomenclature is dynamic. Pango lineage naming and assignment may change as more samples are sequenced and analyzed globally.
- Whole genome sequencing sample logistics are complex and require samples to be transferred across a large network of laboratories. We are unable to verify all eligible samples are sent to the OCGN laboratories for sequencing.

- Data submitted to the OHDP-PHAE from OCGN laboratories have not been independently verified.
- The dates associated with samples submitted by network laboratories vary due to sample logistics and different laboratory information systems. Dates associated with WGS samples were assigned based on a hierarchy: sample collection date > SARS-CoV-2 diagnostic received date > SARS-CoV-2 diagnostic reported date > WGS received date > WGS reported date. Weeks were created to align with surveillance weeks used by the Public Health Agency of Canada for influenza reporting.
- Samples from the same case were linked if they had the same health card number or if they had the same first name, last name, and date of birth. If more than one sample was sequenced for a case, the most recent sample was used. This may shift a case to a more recent week if a subsequent sample was sequenced from the same case. A small proportion of cases may have samples that were not linked due to inconsistencies or data entry errors.
- Results for recent weeks are incomplete as not all sequencing and bioinformatics analyses were complete at the time of data extraction.
- Geographic region was assigned based on OCGN patient postal code. 15.2% of cases had missing/unassigned patient postal code, and 0.1% of cases had out of province postal codes.
- North West region includes Northwestern Health Unit and Thunder Bay District Health Unit; North East region includes Algoma Public Health, North Bay Parry Sound District Health Unit, Porcupine Health Unit, Public Health Sudbury & Districts and Timiskaming Health Unit; Eastern region includes Eastern Ontario Health Unit, Hastings Prince Edward Public Health, Kingston, Frontenac and Lennox & Addington (KFLA) Public Health, Leeds, Grenville & Lanark District Health Unit, Ottawa Public Health and Renfrew County and District Health Unit; Central East region includes Durham Region Health Department, Haliburton, Kawartha, Pine Ridge (HKPR) District Health Unit, Peel Public Health, Peterborough Public Health, Simcoe Muskoka District Health Unit, and York Region Public Health; Toronto region includes Toronto Public Health; South West region includes Chatham-Kent Public Health, Grey Bruce Health Unit, Huron Perth Public Health, Lambton Public Health, Middlesex-London Health Unit, Southwestern Public Health, and Windsor-Essex County Health Unit; Central West region includes Brant County Health Unit, City of Hamilton Public Health Services, Haldimand-Norfolk Health Unit, Halton Region Public Health, Niagara Region Public Health, Region of Waterloo Public Health and Emergency Services, and Wellington-Dufferin-Guelph Public Health.
- For representative surveillance, results may not be representative of Ontario overall. Samples selected include a proportion of eligible samples received by OCGN laboratories according to the whole genome sequencing strategy.
 - Data from the OCGN laboratories cover different time periods: PHO since January 1, 2021, HSC since April 21, 2021, KHSC since January 1, 2021, SHL since March 26, 2021, and HRLMP since April 11, 2021.

Methods: Nowcast Estimates, Projections and Weekly Relative Growth Rate

- Nowcast estimates and projections are generated using a multinomial logistic regression model. The Nowcast model uses six weeks of daily representative surveillance data up to the most recent date, with date as the univariate model predictor. Lineages that had at least one day with an estimated or projected prevalence of 5% or greater were included in the table and figure. Lineages with at least 21 days of non-zero case counts were included in the model.
- Projected Nowcast estimates are future predictions of prevalence after the most recent date of observed data.

- Weekly relative growth rate is a measure of a lineage's growth relative to a reference lineage, over a seven day period.⁶ Relative growth rates greater than one suggest an increased growth rate compared to the reference; relative growth rates less than one suggest a decreased growth rate compared to the reference.
- These weekly relative growth rates can be calculated by exponentiating the weekly selection rate coefficient from the Nowcast model.
 - The selection rate coefficient is the difference in growth rate between two lineages ($\Delta r = r_{\text{lineage}} - r_{\text{reference}}$), and can be derived from a logistic regression model where the outcome is the relative frequency of a lineage and the predictor is the date, measured in weeks.^{6,7}
- The weekly relative growth rate and projections may be overestimated for emerging lineages.

Data Caveats and Methods: Ontario Laboratories Information System Data – Ontario Respiratory Virus Tool

- Sample collection date is used to assign the date of the test.
- The number of tests performed does not reflect the number of specimens or persons tested. More than one test may be performed per specimen or per person. As such, the number of positive tests does not necessarily translate to the number of specimens or persons testing positive. For more information about this data source, see [ORVT Technical Notes](#).

References

1. Public Health Agency of Canada. SARS-CoV-2 variants: national definitions, classifications and public health actions [Internet]. Ottawa, ON: Government of Canada; 2021 [modified 2024 August 22; cited 2021 August 27]. Available from: <https://www.canada.ca/en/public-health/services/diseases/2019-novel-coronavirus-infection/health-professionals/testing-diagnosing-case-reporting/sars-cov-2-variants-national-definitions-classifications-public-health-actions.html>
2. World Health Organization. Tracking SARS-CoV-2 variants [Internet]. Geneva, Switzerland; 2021 [modified 2024 December 9; cited 2021 November 1]. Available from: <https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/>
3. European Centre for Disease Prevention and Control. SARS-CoV-2 variants of concern as of 20 December 2024 [Internet]. Stockholm, Sweden; 2021 [modified 2025 January 3; cited 2021 November 1]. Available from: <https://www.ecdc.europa.eu/en/covid-19/variants-concern>
4. Rambaut A, Holmes EC, O'Toole Á, Hill V, McCrone JT, Ruis C, et al. A dynamic nomenclature proposal for SARS-CoV-2 lineages to assist genomic epidemiology. *Nat Microbiol*. 2020; 5(11):1403-7. Available from: <https://doi.org/10.1038/s41564-020-0770-5>
5. cov-lineages. pangolin [Internet]. 2020 [cited 2021 May 29]. GitHub. Available from: <https://github.com/cov-lineages/pangolin>
6. Davies N, Abbott S, Barnard R, Jarvis CI, Kucharski AJ, Munday JD, et al. Estimated transmissibility and impact of SARS-CoV-2 lineage B.1.1.7 in England. *Science*. 2021; 372(6538): eabg3055. Available from: <https://doi.org/10.1126/science.abg3055>
7. Campbell F, Archer B, Laurenson-Schafer H, Jinnai Y, Konings F, Batra N, et al. Increased transmissibility and global spread of SARS-CoV-2 variants of concern as at June 2021. *Euro Surveill*. 2021; 26(24). Available from: <https://doi.org/10.2807/1560-7917.ES.2021.26.24.2100509>

Citation

Ontario Agency for Health Protection and Promotion (Public Health Ontario). Epidemiologic summary: SARS-CoV-2 whole genome sequencing in Ontario, February 3, 2025. Toronto, ON: King's Printer for Ontario; 2025.

Disclaimer

This document was developed by Public Health Ontario (PHO). PHO provides scientific and technical advice to Ontario's government, public health organizations and health care providers. PHO's work is guided by the current best available evidence at the time of publication. The application and use of this document is the responsibility of the user. PHO assumes no liability resulting from any such application or use. This document may be reproduced without permission for non-commercial purposes only and provided that appropriate credit is given to PHO. No changes and/or modifications may be made to this document without express written permission from PHO.

Public Health Ontario

Public Health Ontario is an agency of the Government of Ontario dedicated to protecting and promoting the health of all Ontarians and reducing inequities in health. Public Health Ontario links public health practitioners, front-line health workers and researchers to the best scientific intelligence and knowledge from around the world.

For more information about PHO, visit publichealthontario.ca.

© King's Printer for Ontario, 2025

Ontario 