





## Vaping and Susceptibility to COVID-19 Infection

No studies were identified that directly measure the relationship between individuals who vape and whether they experience an increase in risk of COVID-19 infection. Thirteen articles were identified that discuss COVID-19 susceptibility among individuals who vape.<sup>1,2,4,6,7,18-26</sup> Vaping products are relatively new; comparisons among studies are challenging because they are evolving in design, diversity of ingredients and functions.<sup>3</sup> Vaping products often contain nicotine and produce aerosol instead of smoke. The use of vaping products has been associated with lung damage.<sup>27,28</sup> Independent of nicotine exposure, particulates and flavourings in e-cigarette/vape aerosols could also potentially impair lung function.<sup>3</sup>

## Vaping Behaviour

Three articles reported vaping behaviour that may be associated with COVID-19 transmission, unrelated to inhalation.<sup>4,5,24</sup> The act of vaping involves frequent hand-to-mouth contact which may increase opportunities to contract COVID-19.<sup>4,5</sup> It is also common for individuals who vape to share devices, and to vape in a group setting and/or confined spaces.<sup>5</sup> These types of actions increase the possibility of viral transmission from hand to mouth and being in close contact with others which are known risk factors for transmission of COVID-19.<sup>4,5,24</sup> Recent data from experimental studies have found that the virus remains stable for several hours to days on surfaces, which makes it plausible for the virus to be transmitted via surfaces of vaping devices.<sup>24</sup>

## COVID-19 Diagnosis Associated With Vaping

Two identified studies analyzed relationships between vaping and COVID-19; one is a population-based analysis using individual-level data on risk factors and the second is an ecological study.<sup>1,2</sup>

Gaiha et. al.<sup>2</sup> conducted a population-based study of youth and young adults to determine whether there was an association between youth smoking, vaping and COVID-19. They found that among young people (ages 13-24 years), a COVID-19 diagnosis was five times more likely among ever users of vaping products only, seven times more likely among ever dual use of vaping products and tobacco cigarettes, and 6.8 times more likely among individuals with dual use in the past 30-days compared to youth who have never vaped before. These findings were adjusted for several major confounders, such as age, sex and obesity. The researchers concluded that COVID-19 is associated with youth use of vaping devices only and dual use of vaping devices and tobacco cigarettes. The authors suggested that their findings have implications for health care providers to screen all youth, especially those that have been infected with COVID-19, for current and a history of vaping. They also highlighted the importance of education in a variety of settings, such as schools, homes, and community-based organizations to support the learning among youth on the use of vaping products, dual use and the impact on the respiratory and immune systems. The authors highlighted the importance of dissemination of youth-focused COVID-19 prevention messaging which would include vaping and dual use.

An ecological study (observational study at the population level) pre-print by Li et. al.<sup>1</sup> indicates some jurisdictional-level correlation in vaping prevalence and proportion of COVID-19 infections and associated deaths in each United States (US) state. Although the study design has limitations and potential confounding effects (e.g., socioeconomic status, underlying health status), the findings do suggest a plausibility of association.

## Potential Role of Cell Receptors

Smoking is one predictor of an individual's likelihood of developing a viral infection, especially a respiratory infection.<sup>19,25</sup> There are hypotheses that individuals who vape would also be at increased risk of developing viral infections. More research is needed to examine these hypotheses.

The angiotensin-converting enzyme 2 (ACE2) is the binding site for the virus causing COVID-19, facilitating its entry into the body.<sup>23</sup> There are studies that have found that nicotine (from smoking tobacco cigarettes) creates an increase in ACE2 expression in the body.<sup>20,21</sup> Given the affinity of COVID-19 for ACE2, these findings raise a concern that nicotine vaping products may also put users at greater risk of COVID-19.<sup>20-22,25</sup> ACE2 levels have also been associated with influencing disease progression.<sup>18,19</sup>

The vast majority of the studies on ACE2 and nicotine have been based on smoking tobacco. While it has been suggested that there would be a similar result among those who use other tobacco and nicotine products, such as vape products, studies are needed to confirm this hypothesis.<sup>18,20-22,25,26</sup>

Further research is needed to determine the role that vaping plays in the susceptibility to and severity of COVID-19 and the relationship with ACE2.

## Vaping, COVID-19 Severity and Outcomes

There is preliminary evidence around the potential role of smoking and/or vaping on COVID-19 infection and severity.<sup>29</sup> Three review articles highlighted indirect studies that reported individuals who vape/smoke may be at a higher risk for severe COVID-19 illness or need mechanical ventilation when compared to non-smokers.<sup>25,30,31</sup> It is hypothesized that the increased risk of contracting COVID-19 and the eventual complications among individuals who smoke/vape are due to delayed clearance of the virus.<sup>25,30</sup> Individuals who vape and smoke tobacco cigarettes (dual users) are considered to be at an increased risk of COVID-19 severe illness since they continue to be exposed to tobacco smoke.<sup>4</sup>

E-cigarette or vaping associated lung injury (EVALI) is an inflammatory response in the lungs triggered by inhaled substances.<sup>16,17</sup> EVALI was newly identified in North America in August 2019. There are concerns of possible coexisting infection with COVID-19 since many of the EVALI symptoms are similar to those of COVID-19.<sup>26,32</sup> Harrill et. al. highlighted concern that in the US, the footprint of the EVALI epidemic nearly mirrors that of current COVID-19 pandemic reported cases, thus matching regional social behavior risk with current regional pandemic risk.<sup>33</sup> EVALI risk among the population who vape represents a possible increased risk of severe COVID-19 illness.<sup>33</sup>

Public health organizations and medical experts around the world are raising concerns that those who smoke and/or vape may be at higher risk for COVID-19 infection susceptibility, severity, and adverse outcomes.<sup>8,34-39</sup> Recently, the CDC reported that adults between 20-44 years of age comprised 20% of COVID-19 hospitalizations in the US.<sup>18,40</sup>

More evidence is needed to determine the impact of vaping on the lungs and whether this impact increases an individual's susceptibility to COVID-19 infection and severity of COVID-19 illness.<sup>26,32</sup>

## Aerosol and COVID-19 Transmission

No studies were identified that directly measure the relationship between aerosol from vaping products and increased risk of COVID-19 infection. However, there is evidence for COVID-19 transmission via

aerosols.<sup>6,23</sup> There is also evidence highlighting the potential for aerosol transmission for other respiratory illnesses such as Middle East Respiratory Syndrome-associated coronavirus (MERS-CoV), Ebola virus<sup>41</sup> and influenza.<sup>42</sup>

The predominant mode of transmission of COVID-19 is via respiratory droplets during close, unprotected contact.<sup>23</sup> Aerosols may be generated during aerosol generating medical procedures (AGMPs), which may increase the risk of transmission.<sup>23</sup> Therefore, it is plausible that the aerosols generated from vaping products may increase COVID-19 transmission.

The WHO acknowledges that there is no evidence about the relationship between vaping and COVID-19, they suggest that given COVID-19 affects the respiratory tract, the hand-to-mouth action of vaping may increase the risk of infection.<sup>7</sup> Additionally, the BC Centre for Disease Control states that while we do not know if one can contract COVID-19 from e-cigarette aerosol, they recommend staying two metres away from others and vaping outside to reduce the risk of infection. They also suggest, that due to the higher risk of complications of COVID-19 among individuals who vape, they recommend reducing or quitting vaping to reduce the risk.<sup>8</sup> This is concerning as the prevalence of e-cigarette use among Canadian youth has increased substantially between 2013 and 2019.<sup>43</sup>

## References

1. Li D, Croft DP, Ossip DJ, Xie Z. Are vapers more susceptible to COVID-19 infection? MedRxiv 20092379 [Preprint]. 2020 May 09 [cited 2020 Sep 15]. Available from: <https://dx.doi.org/10.1101/2020.05.05.20092379>
2. Gaiha SM, Cheng J, Halpern-Felsher B. Association between youth smoking, electronic cigarette use, and coronavirus disease 2019. *J Adolesc Health*. 2020;67(4):519-23. Available from: <https://doi.org/10.1016/j.jadohealth.2020.07.002>
3. National Academies of Sciences, Engineering, and Medicine. Public health consequences of e-cigarettes. Washington, DC: National Academies Press; 2018. Available from: <https://doi.org/10.17226/24952>
4. Ontario Agency for Health Protection and Promotion (Public Health Ontario). Tobacco and COVID-19 – what we know so far [Internet]. Toronto, ON: Queen’s Printer for Ontario; 2020 [cited 2020 Sep 30]. Available from: <https://www.publichealthontario.ca/-/media/documents/nCoV/covid-wwksf/2020/08/what-we-know-tobacco-covid.pdf?la=en>
5. Majmundar A, Allem JP, Cruz TB, Unger JB. Public health concerns and unsubstantiated claims at the intersection of vaping and COVID-19. *Nicotine Tobacco Res*. 2020;22(9):1667-8. Available from: <https://doi.org/10.1093/ntr/ntaa064>
6. Ontario Agency for Health Protection and Promotion (Public Health Ontario). Review of “It is time to address airborne transmission of COVID-19” [Internet]. Toronto, ON: Queen's Printer for Ontario; 2020 [cited 2020 Sep 30]. Available from: <https://www.publichealthontario.ca/-/media/documents/nCoV/research/2020/07/research-morawska-clinfectdis-its-time-to-address-airborne-transmission.pdf?la=en>
7. World Health Organization. Q&A: tobacco and COVID-19 [Internet]. Geneva: World Health Organization; 2020 [modified May 27; cited 2020 Sep 15]. Available from:

<https://www.who.int/emergencies/diseases/novel-coronavirus-2019/question-and-answers-hub/q-a-detail/q-a-on-smoking-and-covid-19#>

8. BC Centre for Disease Control. Vaping, smoking and COVID-19 [Internet]. Vancouver, BC: Provincial Health Services Authority; 2020 [cited 2020 Sep 04]. Available from: <http://www.bccdc.ca/health-info/diseases-conditions/covid-19/prevention-risks/vaping-smoking-and-covid-19>

9. Ontario Agency for Health Protection and Promotion (Public Health Ontario), Berenbaum E, Keller-Olaman S, Manson H, Moloughney B, Muir S, et al. Current evidence on e-cigarettes: a summary of potential impacts [Internet]. Toronto, ON: Queen's Printer for Ontario; 2018 [cited 2020 Sep 30]. Available from: <https://www.publichealthontario.ca/-/media/documents/l/2018/literature-review-ecigarettes.pdf?la=en>

10. Statistics Canada. Canadian tobacco and nicotine survey, 2019 [Internet]. Ottawa, ON: Government of Canada; 2020 [modified 2020 Mar 05; cited 2020 Sep 21]. Available from: <https://www150.statcan.gc.ca/n1/daily-quotidien/200305/dq200305a-eng.htm>

11. Ontario Agency for Health Protection and Promotion (Public Health Ontario). Ontario tobacco monitoring report 2018. Toronto, ON: Queen's Printer for Ontario; 2019. Available from: <https://www.publichealthontario.ca/-/media/documents/t/2019/tobacco-report-2018.pdf?la=en>

12. Han L, Ran J, Mak YW, Suen LK, Lee PH, Peiris JSM, et al. Smoking and influenza-associated morbidity and mortality: a systematic review and meta-analysis. *Epidemiology*. 2019;30(3):405-17. Available from: <https://doi.org/10.1097/ede.0000000000000984>

13. U.S. Department of Health and Human Services, Public Health Service, Office of the Surgeon General. The health consequences of smoking—50 years of progress: a report of the Surgeon General [Internet]. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2014 [cited 2020 Sep 30]. Available from: [https://www.cdc.gov/tobacco/data\\_statistics/sgr/50th-anniversary/index.htm](https://www.cdc.gov/tobacco/data_statistics/sgr/50th-anniversary/index.htm)

14. Pushalkar S, Paul B, Li Q, Yang J, Vasconcelos R, Makwana S, et al. Electronic cigarette aerosol modulates the oral microbiome and increases risk of infection. *iScience*. 2020;23(3):100884. Available from: <https://doi.org/10.1016/j.isci.2020.100884>

15. Gotts JE, Jordt S, McConnell R, Tarran R. What are the respiratory effects of e-cigarettes? *BMJ*. 2019;366:l5275. Available from: <https://doi.org/10.1136/bmj.l5275>

16. Government of Canada. Vaping-associated lung illness [Internet]. Ottawa, ON: Government of Canada; 2020 [modified 2020 Aug 20; cited 2020 Sep 04]. Available from: <https://www.canada.ca/en/public-health/services/diseases/vaping-pulmonary-illness.html>

17. Centers for Disease Control and Prevention. Outbreak of lung injury associated with the use of e-cigarette, or vaping, products [Internet]. Atlanta, GA: Centers for Disease Control and Prevention; 2020 [modified 2020 Feb 25; cited 2020 Sep 04]. Available from: [https://www.cdc.gov/tobacco/basic\\_information/e-cigarettes/severe-lung-disease.html](https://www.cdc.gov/tobacco/basic_information/e-cigarettes/severe-lung-disease.html)

18. Pino LE, Triana I, Perez C, Piotrostanalzki A, Ruiz-Patino A, Lopes G, et al. Electronic nicotine delivery systems (ECs) and COVID-19: the perfect storm for young consumers. *Clin Transl Oncol*. 2020 May 23 [Epub ahead of print]. Available from: <https://doi.org/10.1007/s12094-020-02391-x>
19. Kabbani N, Olds JL. Does COVID19 infect the brain? If so, smokers might be at a higher risk. *Mol Pharmacol*. 2020;97(5):351-3. Available from: <https://doi.org/10.1124/molpharm.120.000014>
20. Sharma P, Zeki AA. Does vaping increase susceptibility to COVID-19? *Am J Respir Crit Care Med*. 2020 Aug 04 [Epub ahead of print]. Available from: <https://doi.org/10.1164/rccm.202005-2103le>
21. McAlinden KD, Eapen MS, Lu W, Chia C, Haug G, Sohal SS. COVID-19 and vaping: risk for increased susceptibility to SARS-CoV-2 infection? *Eur Respir J*. 2020;56(1):2001645. Available from: <https://doi.org/10.1183/13993003.01645-2020>
22. Lee AC, Chakladar J, Li WT, Chen C, Chang EY, Wang-Rodriguez J, et al. Tobacco, but not nicotine and flavor-less electronic cigarettes, induces ACE2 and immune dysregulation. *Int J Mol Sci*. 2020;21(15):5513. Available from: <https://doi.org/10.3390/ijms21155513>
23. Ontario Agency for Health Protection and Promotion (Public Health Ontario). COVID-19 routes of transmission – what we know so far [Internet]. Toronto, ON: Queen's Printer for Ontario; 2020 [cited 2020 Sep 29]. Available from: <https://www.publichealthontario.ca/-/media/documents/ncov/wwksf-routes-transmission-mar-06-2020.pdf?la=en>
24. Kampf G, Todt D, Pfaender S, Steinmann E. Persistence of coronaviruses on inanimate surfaces and their inactivation with biocidal agents. *J Hosp Infect*. 2020;104(3):246-51. Available from: <https://doi.org/10.1016/j.jhin.2020.01.022>
25. Kaur G, Lungarella G, Rahman I. SARS-CoV-2 COVID-19 susceptibility and lung inflammatory storm by smoking and vaping. *J Inflamm (Lond)*. 2020;17:21. Available from: <https://doi.org/10.1186/s12950-020-00250-8>
26. *Lancet Respir Med*. The EVALI outbreak and vaping in the COVID-19 era. *Lancet Respir Med*. 2020;8(9):831. Available from: [https://doi.org/10.1016/s2213-2600\(20\)30360-x](https://doi.org/10.1016/s2213-2600(20)30360-x)
27. Brake SJ, Barnsley K, Lu W, McAlinden KD, Eapen MS, Sohal SS. Smoking upregulates angiotensin-converting enzyme-2 receptor: a potential adhesion site for novel coronavirus SARS-CoV-2 (covid-19). *J Clin Med*. 2020;9(3):841. Available from: <https://doi.org/10.3390/jcm9030841>
28. Javelle E. Electronic cigarette and vaping should be discouraged during the new coronavirus SARS-CoV-2 pandemic. *Arch Toxicol*. 2020;94(6):2261-2. Available from: <https://dx.doi.org/10.1007%2Fs00204-020-02744-z>
29. Russo P, Bonassi S, Giacconi R, Malavolta M, Tomino C, Maggi F. COVID-19 and smoking: is nicotine the hidden link? *Eur Respir J*. 2020;55(6):2001116. Available from: <https://doi.org/10.1183/13993003.01116-2020>
30. Munzel T, Hahad O, Kuntic M, Keaney JF, Deanfield JE, Daiber A. Effects of tobacco cigarettes, e-cigarettes, and waterpipe smoking on endothelial function and clinical outcomes. *Eur Heart J*. 2020 Jun 25 [Epub ahead of print]. Available from: <https://dx.doi.org/10.1093/eurheartj/ehaa460>

31. Archie SR, Cucullo L. Cerebrovascular and neurological dysfunction under the threat of COVID-19: is there a comorbid role for smoking and vaping? *Int J Mol Sci.* 2020;21(11):3916. Available from: <https://dx.doi.org/10.3390/ijms21113916>
32. Galo J, Celli D, Gross D, Holt G, Campos M. A presentation of e-cigarette vaping associated lung injury (EVALI) caused by THC-containing electronic smoking device. *Respir Med Case Rep.* 2020;31:101154. Available from: <https://doi.org/10.1016/j.rmcr.2020.101154>
33. Harrill WC. Vaping during the COVID-19 pandemic: NOT GOOD!! *Laryngoscope Investig Otolaryngol.* 2020;5(3):399-400. Available from: <https://doi.org/10.1002/lio2.386>
34. World Health Organization Regional Office for the Eastern Mediterranean. Tobacco free initiative: tobacco and waterpipe use increases the risk of COVID-19 [Internet]. Geneva: World Health Organization; 2020 [cited 2020 Sep 29]. Available from: <http://www.emro.who.int/tfi/know-the-truth/tobacco-and-waterpipe-users-are-at-increased-risk-of-covid-19-infection.html>
35. Chadi N, Bélanger R. COVID, youth, and substance use: critical messages for youth and families [Internet]. Ottawa, ON: Canadian Paediatric Society; 2020 [modified 2020 Apr 20; cited 2020 Sep 21]. Available from: <https://www.cps.ca/en/blog-blogue/covid-youth-and-substance-use-critical-messages-for-youth-and-families>
36. Centers for Disease Control and Prevention. People with certain medical conditions [Internet]. Atlanta, GA: Centers for Disease Control and Prevention; 2020 [modified 2020 Sep 11; cited 2020 Sep 21]. Available from: [https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/people-with-medical-conditions.html?CDC\\_AA\\_refVal=https%3A%2F%2Fwww.cdc.gov%2Fcoronavirus%2F2019-ncov%2Fneed-extra-precautions%2Fgroups-at-higher-risk.html](https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/people-with-medical-conditions.html?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Fcoronavirus%2F2019-ncov%2Fneed-extra-precautions%2Fgroups-at-higher-risk.html)
37. Public Health Agency of Canada. Statement from the Chief Public Health Officer of Canada on COVID-19 [Internet]. Ottawa, ON: Government of Canada [modified 2020 May 31; cited 2020 Sep 21]. Available from: <https://www.canada.ca/en/public-health/news/2020/05/statement-from-the-chief-public-health-officer-of-canada-on-covid-1910.html>
38. Australian Government, Department of Health. Australian Health Protection Principal Committee (AHPPC) statement on tobacco use, e-cigarette use and COVID-19 [Internet]. Canberra: Commonwealth of Australia; 2020 [cited 2020 Sep 21]. Available from: <https://www.health.gov.au/news/australian-health-protection-principal-committee-ahppc-statement-on-tobacco-use-e-cigarette-use-and-covid-19>
39. Centre for Addiction and Mental Health (CAMH). Vaping and COVID-19: information for people who use e-cigarettes [Internet]. Toronto, ON: CAMH; 2020 [cited 2020 Sep 21]. Available from: [https://camh.ca/-/media/files/camh\\_covid19\\_infosheet-ndc-vaping-pdf.pdf](https://camh.ca/-/media/files/camh_covid19_infosheet-ndc-vaping-pdf.pdf)
40. CDC COVID-19 Response Team. Severe outcomes among patients with coronavirus disease 2019 (COVID-19) - United States, February 12-March 16, 2020. *MMWR Morb Mortal Wkly Rep.* 2020;69(12):343-6. Available from: <https://doi.org/10.15585/mmwr.mm6912e2>
41. Tellier R, Li Y, Cowling BJ, Tang JW. Recognition of aerosol transmission of infectious agents: a commentary. *BMC Infect Dis.* 2019;19(1):101. Available from: <https://doi.org/10.1186/s12879-019-3707-y>



42. Yan J, Grantham M, Pantelic J, Bueno de Mesquita PJ, Albert B, Liu F, et al. Infectious virus in exhaled breath of symptomatic seasonal influenza cases from a college community. *Proc Natl Acad Sci U S A*. 2018;115(5):1081-6. Available from: <https://doi.org/10.1073/pnas.1716561115>

43. Cole AG, Aleyan S, Battista K, Leatherdale ST. Trends in youth e-cigarette and cigarette use between 2013 and 2019: insights from repeat cross-sectional data from the COMPASS study. *Can J Public Health*. 2020 Aug 17 [Epub ahead of print]. Available from: <https://doi.org/10.17269/s41997-020-00389-0>

## Citation

Ontario Agency for Health Protection and Promotion (Public Health Ontario). Vaping and COVID-19 – what we know so far. Toronto, ON: Queen’s Printer for Ontario; 2020.

## Disclaimer

This document was developed by Public Health Ontario (PHO). PHO provides scientific and technical advice to Ontario’s government, public health organizations and health care providers. PHO’s work is guided by the current best available evidence at the time of publication.

The application and use of this document is the responsibility of the user. PHO assumes no liability resulting from any such application or use.

This document may be reproduced without permission for non-commercial purposes only and provided that appropriate credit is given to PHO. No changes and/or modifications may be made to this document without express written permission from PHO.

## Public Health Ontario

Public Health Ontario is an agency of the Government of Ontario dedicated to protecting and promoting the health of all Ontarians and reducing inequities in health. Public Health Ontario links public health practitioners, front-line health workers and researchers to the best scientific intelligence and knowledge from around the world.

For more information about PHO, visit [publichealthontario.ca](https://publichealthontario.ca).

